
Burns Hesi Case Study

*Downloaded from
remodelicious.com by
Herman & Jaiden*

Burns Hesi Case Study

INTRODUCTION: MASTERING THE BURNS HESI CASE STUDY

Nursing students often encounter few clinical scenarios as demanding and multifaceted as the care of a burn-injured patient. The **Burns Hesi Case Study** is a staple of the HESI specialty exams, designed to test not only your recall of pathophysiology but also your ability to prioritize care, interpret complex clinical data, and make rapid

critical decisions. A burn injury affects virtually every organ system, from the integumentary and respiratory systems to fluid balance, immune function, and psychosocial well-being. Mastering this type of case study requires a deep understanding of the sequential phases of burn management, the nuanced calculations for fluid resuscitation, and the subtle signs of complications such as inhalation injury or sepsis. This article provides a thorough exploration of the Burns HESI case study, offering practical frameworks, evidence-based insights, and study strategies to help you excel

on the exam and, more importantly, in clinical practice.

UNDERSTANDING THE SCOPE OF BURN INJURIES IN HESI CASE STUDIES

Before diving into specific case details, it is essential to grasp why burn injuries are a favorite topic for HESI exam writers. Burn cases uniquely integrate content from medical-surgical nursing, pharmacology, nutrition, mental health, and even pediatric or geriatric nursing. A typical **Burns Hesi Case Study** will present a patient with a specific percentage of total body surface area (TBSA) burned, a certain depth of injury, and often a complicating factor such as inhalation injury, pre-existing comorbidities, or age extremes. The

exam expects you to identify the immediate priorities, calculate fluid requirements using the Parkland formula, anticipate the patient's trajectory through the emergent, acute, and rehabilitation phases, and recognize complications like Curling's ulcer or compartment syndrome.

Key Components You Will Encounter

When you open a Burns HESI case study, you are typically given a brief narrative. For example, a 45-year-old male is brought to the emergency department

after a house fire, with burns to his anterior chest, both arms, and his face. The case will provide vital signs, weight, and possibly initial laboratory values. From this, you must determine the TBSA burned using the Rule of Nines, classify the burn depth (superficial partial-thickness, deep partial-thickness, or full-thickness), and then prioritize interventions. The case study will progress through time, often presenting changes in the patient's condition—such as decreased urine output, rising potassium levels, or signs of respiratory distress—requiring you to adjust your plan of care accordingly.

THE PATHOPHYSIOLOGY FOUNDATION: WHY BURN CARE

IS UNIQUE

To answer questions accurately in a **Burns Hesi Case Study**, you need a solid grasp of the pathophysiology. A severe burn destroys the skin's protective barrier, leading to massive fluid shifts, heat loss, and a portal for infection. The systemic inflammatory response syndrome (SIRS) triggered by the burn injury can cause capillary leak, hypovolemia, and, eventually, distributive shock. Remember that the initial 24 to 48 hours are dominated by hypovolemia and the need for aggressive fluid resuscitation. After the capillary leak seals, the patient enters a hypermetabolic, catabolic state that can last for weeks or months. This phase is characterized by increased metabolic

rate, protein wasting, hyperglycemia, and immunosuppression. Understanding this biphasic response is critical for predicting the patient's needs and prioritizing nursing interventions.

Inhalation Injury: The Silent Threat

Many burn cases involve inhalation injury, which dramatically increases morbidity and mortality. In a Burns HESI case study, always look for clues such as burns on the face, singed nasal hairs, carbonaceous sputum, hoarseness, or a history of being trapped in an enclosed space. Inhalation injury can cause upper airway edema, lower airway damage, and carbon monoxide poisoning. The

priority here is airway management. These patients often require early intubation because airway swelling can progress rapidly. Pulse oximetry may be falsely elevated in carbon monoxide poisoning, so a carboxyhemoglobin level is a more reliable indicator. The key nursing action is to administer 100% humidified oxygen until the carboxyhemoglobin level falls below 10%.

FLUID RESUSCITATION IN THE BURNS HESI CASE STUDY

One of the most frequently tested topics is the Parkland formula for fluid resuscitation. In any **Burns Hesi Case Study**, you will likely need to calculate the patient's fluid requirements for the first 24 hours. The formula is 4 mL of

Lactated Ringer's solution per kilogram of body weight per percentage of TBSA burned. Half of this total is given in the first 8 hours from the time of injury, and the remaining half is given over the next 16 hours. It is crucial to use the time of injury, not the time of arrival to the hospital, as the starting point. HESI questions often test your ability to adjust the rate of infusion based on urine output, which should be maintained at 30 to 50 mL per hour in adults (or 1 mL/kg/hour in children). If urine output falls below this threshold, the nurse should increase the IV rate; if it exceeds 75 mL/hour, the rate should be decreased to prevent fluid overload.

Monitoring Endpoints of Resuscitation

Beyond urine output, the Burns HESI case study may require you to monitor other endpoints of resuscitation, including heart rate, blood pressure, central venous pressure, and acid-base status. Lactate levels and base deficit are valuable indicators of tissue perfusion. A rising lactate or worsening base deficit suggests inadequate resuscitation. However, remember that fluid overload can lead to abdominal compartment syndrome or pulmonary edema. The nurse must balance

between under-resuscitation, which leads to acute kidney injury, and over-resuscitation, which can cause respiratory compromise. This delicate balance is a hallmark of expert burn care.

WOUND CARE AND INFECTION PREVENTION

After the initial resuscitation phase, the focus shifts to wound care and infection prevention. In a **Burns Hesi Case Study**, you may be asked about the appropriate topical antimicrobial agents for different depths of burns. Silver sulfadiazine is commonly used for partial-thickness burns, but it is contraindicated on the face due to staining. Mafenide acetate penetrates eschar well and is effective against

gram-negative organisms, but it is painful and can cause metabolic acidosis. Bacitracin or bacitracin-polymyxin B ophthalmic ointment is used for facial burns, especially around the eyes. The case study may also ask about the appropriate dressing technique, the use of enzymatic debriding agents like collagenase, or the signs of wound infection, such as a change in exudate color, foul odor, periwound erythema, or an increase in pain.

Escharotomy and

Fasciotomy

Full-thickness burns create an inelastic eschar that can restrict chest expansion or compromise circulation to a limb. The Burns HESI case study often includes a scenario where the patient develops signs of compartment syndrome, such as deep pain, pallor, paresthesia, pulselessness, or paralysis. In circumferential full-thickness burns of the chest, escharotomy is performed to allow for adequate chest wall excursion and ventilation. In circumferential burns of an extremity, escharotomy or fasciotomy is performed to restore distal circulation. The nurse's role includes frequent neurovascular assessments, monitoring for pain out of proportion to

the injury, and preparing the patient for the procedure.

NUTRITIONAL AND METABOLIC SUPPORT

Burn injury induces a profound hypermetabolic state that can increase resting energy expenditure by 100% or more. A significant portion of the **Burns Hesi Case Study** may focus on nutritional management. Early enteral nutrition, ideally within 24 to 36 hours of injury, is the standard of care to maintain gut barrier integrity and reduce the risk of stress ulcers. The diet should be high in protein and calories. The case study may ask you to calculate the patient's caloric needs using the Curreri formula or to identify the need for vitamin and mineral supplementation,

particularly vitamin A, vitamin C, zinc, and iron. Monitoring for complications such as feeding intolerance, diarrhea, or hyperglycemia is an essential nursing responsibility.

Stress Ulcer Prophylaxis

Curling's ulcer, a stress ulcer that occurs in the duodenum of burn patients, was once a common complication. Today, routine prophylaxis with histamine H2-receptor blockers or proton pump inhibitors has significantly reduced its incidence. In a Burns HESI case study, you may need to recognize the signs of gastrointestinal bleeding, such as coffee-ground emesis or melena, and identify

the appropriate prophylactic medication.

PSYCHOSOCIAL AND EMOTIONAL IMPACT

Burn injuries are devastating not only physically but also emotionally and socially. The Burns HESI case study will often include a mental health component, assessing the patient's coping mechanisms, body image disturbance, and risk for post-traumatic stress disorder (PTSD). Pain management is closely tied to psychosocial care, as unrelieved pain exacerbates anxiety and depression. The case study may ask about the appropriate use of patient-controlled analgesia (PCA), the role of anxiolytics, and the importance of nonpharmacologic interventions such as guided imagery,

relaxation techniques, or virtual reality therapy. Additionally, the case may involve the family's response to the injury and the need for social work or chaplaincy referral.

Pediatric and Geriatric Considerations

Age extremes are frequently featured in Burns HESI case studies. In children, the Rule of Nines is adjusted because their body proportions are different. The head accounts for a larger percentage (18% in children vs. 9% in adults), and the legs account for a smaller proportion.

Children also have a higher BSA-to-mass ratio, making them more prone to hypothermia and hypovolemia. In older adults, the skin is thinner, healing is slower, and pre-existing conditions such as diabetes, hypertension, or coronary artery disease complicate care. The case study may test your ability to modify the Parkland formula or adjust medication doses based on age-related changes in renal or hepatic function.

STUDY STRATEGIES FOR THE BURNS HESI CASE STUDY

To succeed on the Burns HESI case study, you must move beyond memorizing facts and develop a systematic approach to clinical reasoning. Here are some practical strategies.

Use the Nursing Process

Organize your thinking using the steps of the nursing process: assessment, diagnosis, planning, implementation, and evaluation. When you read the case study, actively identify the data that supports each step. For example, if the patient has full-thickness burns on both legs and decreased pedal pulses, your nursing diagnosis might be "Ineffective peripheral tissue perfusion related to edema." Your intervention might be to elevate the extremities and prepare for escharotomy, and your evaluation criteria would be the return of palpable pulses.

Prioritize Airway, Breathing, Circulation

Burn cases can be overwhelming with the amount of data presented. Always start with the ABCs. Is the airway patent? Are there signs of inhalation injury? Is the patient breathing adequately? Is there circumferential chest burn restricting expansion? What is the heart rate and blood pressure? What is the urine output? By anchoring yourself to these foundational priorities, you can avoid getting lost in less critical details.

Master the Calculations

The Parkland formula is a must-know. Practice several variations so that you can compute it quickly and accurately. Also, be comfortable with the Rule of Nines for adults and the modified version for children. HESI questions often require you to calculate the TBSA from a description of the burned areas, so practice identifying which body parts correspond to which percentages.

Review Common

Medications

Be familiar with the indications, dosages, side effects, and nursing considerations for the medications used in burn care, including topical antimicrobials, opioids, anxiolytics, proton pump inhibitors, and insulin. The Burns HESI case study may ask you to identify the correct medication for a specific wound type or the appropriate treatment for hyperkalemia following massive tissue destruction.

Understand the

Phases of Burn Care

The emergent (resuscitative) phase lasts from the time of injury to about 48 hours, focusing on airway management, fluid resuscitation, and prevention of shock. The acute phase begins after successful resuscitation and continues until wound closure, emphasizing wound care, infection prevention, and nutritional support. The rehabilitation phase focuses on scar management, physical and occupational therapy, and psychosocial reintegration. Knowing which phase the patient is in helps you

predict the appropriate interventions and anticipate complications.

PUTTING IT ALL TOGETHER: A PRACTICAL APPLICATION

Imagine you are presented with a Burns HESI case study of a 30-year-old female with 35% TBSA deep partial-thickness burns to her anterior trunk and both arms. She weighs 70 kg. She has stridor and carbonaceous sputum. The time of injury was 21:00, and it is now 02:00. Your first priority is to secure the airway due to suspected inhalation injury. You calculate the 24-hour fluid requirement: $4 \text{ mL} \times 70 \text{ kg} \times 35\% = 9,800 \text{ mL}$. Half of this (4,900 mL) must be given in the first 8 hours from